

Rotman assisted in the preparation and interpretation of thin sections. This work was partly funded by NASA Planetary Geology and Geophysics grant NNG05GJ91G and by the Rhodes Trust.

References

- Arvidson, R. E., et al. (2006), Overview of the Spirit Mars Exploration Rover Mission to Gusev Crater: Landing site to Backstay Rock in the Columbia Hills, *J. Geophys. Res.*, *111*, E02S01, doi:10.1029/2005JE002499.
- Aydin, A., and J. M. Degraff (1988), Evolution of polygonal fracture patterns in lava flows, *Science*, *239*, 471–476, doi:10.1126/science.239.4839.471.
- Baker, V. (1978a), Quaternary geology of the Channeled Scabland and adjacent areas, in *The Channeled Scabland: A Guide to the Geomorphology of the Columbia Basin, Washington*, edited by V. Baker and D. Nummedal, pp. 59–79, NASA, Washington, D. C.
- Baker, V. (1978b), Large-scale erosional and depositional features of the channeled scabland, in *The Channeled Scabland: A Guide to the Geomorphology of the Columbia Basin, Washington*, edited by V. Baker and D. Nummedal, pp. 81–116, NASA, Washington, D. C.
- Baker, V., and D. Nummedal (Eds.) (1978), *The Channeled Scabland: A Guide to the Geomorphology of the Columbia Basin, Washington*, 186 pp., NASA, Washington, D. C.
- Barrett, P. J. (1980), The shape of rock particles: A critical review, *Sedimentology*, *27*, 291–303, doi:10.1111/j.1365-3091.1980.tb01179.x.
- Basilevsky, A. T., W. J. Markiewicz, N. Thomas, and H. U. Keller (1999), Morphologies of rocks within and near the Rock Garden at the Mars Pathfinder landing site, *J. Geophys. Res.*, *104*(E4), 8617–8636, doi:10.1029/1998JE900039.
- Birkeland, P. W. (1999), *Soils and Geomorphology*, 448 pp., Oxford Univ. Press, New York.
- Boelhouwers, J., A. De Joode, and D. Jager (1999), Relative-age dating of debris flow deposits in the Cederberg, *S. Afr. Geogr. J.*, *81*, 135–142.
- Bolongaro-Crevenna, A., V. Torres-Rodriguez, V. Sorani, D. Frame, and M. A. Ortiz (2005), Geomorphometric analysis for characterizing landforms in Morelos State, Mexico, *Geomorphology*, *67*(3–4), 407–422, doi:10.1016/j.geomorph.2004.11.007.
- Bourke, M., H. Viles, J. Nicoli, P. Lyew-Ayee, R. Ghent, and J. Holmlund (2008), Innovative applications of laser scanning and rapid prototype printing to rock breakdown experiments, *Earth Surface Processes Landforms*, doi:10.1002/esp.1631, in press.
- Bourke, M. C., and H. A. Viles (Eds.) (2007), *A Photographic Atlas of Rock Breakdown Features in Geomorphic Environments*, 81 pp., Planet. Sci. Inst., Tucson, Ariz.
- Bretz, J. H. (1923), The Channeled Scabland of the Columbia Plateau, *J. Geol.*, *31*, 617–649.
- Buffin-Bélanger, T., I. Reid, S. Rice, J. H. Chandler, and J. Lancaster (2003), A casting procedure for reproducing coarse-grained sedimentary surfaces, *Earth Planet. Sci. Lett.*, *28*(7), 787–796.
- Clague, J. J., R. Barendregt, R. J. Enkin, and F. Foit Jr. (2003), Paleomagnetic and tephra evidence for tens of Missoula floods in southern Washington, *Geology*, *31*(3), 247–250, doi:10.1130/0091-7613(2003)031<0247:PATEFT>2.0.CO;2.
- Curran, J., B. Smith, and P. Warke (2002), Weathering of igneous rocks during shallow burial in an upland peat environment: observations from the Bronze Age Copney Stone Circle Complex, Northern Ireland, *Catena*, *49*, 139–155, doi:10.1016/S0341-8162(02)00023-1.
- Day, M. J., and A. S. Goudie (1977), Field assessment of rock hardness using the Schmidt test hammer, *Tech. Bull. Br. Geomorphol. Res. Group*, *18*, 19–29.
- Dincer, I., A. Acar, I. Cobano, and Y. Uras (2004), Correlation between Schmidt hardness, uniaxial compressive strength, and Young's modulus for andesites, basalts, and tuffs, *Bull. Eng. Geol. Environ.*, *63*(2), 141–148, doi:10.1007/s10064-004-0230-0.
- Dowman, E. A. (1970), *Conservation in Field Archaeology*, 170 pp., Methuen, London.
- Durian, D. J., H. Bidead, P. Durringer, A. Schroder, and C. M. Marques (2006), What is in a pebble shape?, *Phys. Rev. Lett.*, *97*, 028001, doi:10.1103/PhysRevLett.97.028001.
- Evans, I. S., and C. J. McClean (1995), The land surface is not unifractal: Variograms, cirque scale and allometry, *Z. Geomorphol. Suppl.*, *101*, 127–147.
- Fisher, P., J. Wood, and T. Cheng (2004), Where is Helvellyn? Fuzziness of multi-scale landscape morphometry, *Trans. Inst. Br. Geogr.*, *29*, 106–128, doi:10.1111/j.0020-2754.2004.00117.x.
- Golombek, M., and D. Rapp (1997), Size-frequency distributions of rocks on Mars and Earth analog sites: Implications for future landed missions, *J. Geophys. Res.*, *102*(E2), 4117–4130, doi:10.1029/96JE03319.
- Gordon, S. I., and R. I. Dorn (2005), In-situ weathering rind erosion, *Geomorphology*, *67*(1–2), 97–113, doi:10.1016/j.geomorph.2004.06.011.
- Graham, D. J., and N. G. Midgley (2000), Graphical representation of particle shape using triangular diagrams: An Excel spreadsheet method, *Earth Surf. Processes Landforms*, *25*(13), 1473–1477, doi:10.1002/1096-9837(200012)25:13<1473::AID-ESP158>3.0.CO;2-C.
- Greeley, R., N. T. Bridges, R. O. Kuzmin, and J. E. Laity (2002), Terrestrial analogs to wind-related features at the Viking and Pathfinder landing sites on Mars, *J. Geophys. Res.*, *107*(E1), 5005, doi:10.1029/2000JE001481.
- Grolier, M. J., and J. W. Bingham (1971), Geologic map and sections of parts of Grant, Adams, and Franklin Counties, Washington, *Misc. Geol. Invest. Map I-589*, scale 1:62,500, U.S. Geol. Surv., Dep. of Inter., Washington, D. C.
- Hallet, B., J. S. Walder, and C. W. Stubbs (1991), Weathering by segregation ice growth in microcracks at sustained sub-zero temperatures: Verification from an experimental study using acoustic emissions, *Permafrost Periglacial Processes*, *2*, 283.
- Heslop, E. E. M., H. A. Viles, and M. C. Bourke (2004), Understanding rock breakdown on Earth and Mars: Geomorphological concepts and facet mapping methods, *Lunar Planet. Sci. [CD-ROM]*, XXXV, abstract 1445.
- Hooper, P. R. (2000), Chemical discrimination of Columbia River basalt flows, *Geochem. Geophys. Geosyst.*, *1*, doi:10.1029/2000GC000040.
- Hooper, P. R., V. Camp, S. Reidel, and M. Ross (2007), The origin of the Columbia River flood basalt province: Plume versus nonplume models, in *Plates, Plumes, and Planetary Processes*, edited by G. R. Foulger and D. M. Jurdy, *Geol. Soc. of Am. Spec. Pap. 430*, Geol. Soc. of Am., Boulder, Colo.
- Lancaster, N. (1995), *Geomorphology of Desert Dunes*, 320 pp., Routledge, New York.
- Long, P. E., and B. J. Wood (1986), Structures, textures, and cooling histories of Columbia River basalt flows, *Geol. Soc. Am. Bull.*, *97*, 1144–1155, doi:10.1130/0016-7606(1986)97<1144:STACHO>2.0.CO;2.
- Lyle, P. (2000), The eruption environment of multi-tiered columnar basalt lava flows, *J. Geol. Soc. London*, *157*, 715–722.
- Mahaney, W. C. (2002), *Atlas of Sand Grain Surface Textures and Applications*, 237 pp., Oxford Univ. Press, Oxford, U. K.
- McCarroll, D. (1991), The Schmidt hammer, weathering, and rock surface roughness, *Earth Surf. Processes Landforms*, *16*(5), 477–480, doi:10.1002/esp.3290160510.
- McCarroll, D., and A. Nesje (1996), Rock surface roughness as an indicator of degree of rock surface weathering, *Earth Surf. Processes Landforms*, *21*(10), 963–977, doi:10.1002/(SICI)1096-9837(199610)21:10<963::AID-ESP643>3.0.CO;2-J.
- McClean, C. J., and I. S. Evans (2000), Apparent fractal dimensions from continental scale digital elevation models using variogram methods, *Trans. GIS*, *4*, 361–378, doi:10.1111/1467-9671.00061.
- McFadden, L. D., M. C. Eppes, A. R. Gillespie, and B. Hallet (2005), Physical weathering in arid landscapes due to diurnal variation in the direction of solar heating, *Geol. Soc. Am. Bull.*, *117*(1/2), 161–173, doi:10.1130/B25508.1.
- McGreevy, J. P. (1982), Hydrothermal alteration and earth surface rock weathering: A basalt example, *Earth Surf. Processes Landforms*, *7*, 189–195, doi:10.1002/esp.3290070211.
- Mehring, P. J., Jr. (1985), Late-Quaternary pollen records from the interior Pacific Northwest and northern Great Basin of the United States, in *Pollen Records of Late-Quaternary North American Sediments*, edited by V. M. Bryant Jr. and R. G. Holloway, pp. 167–187, Am. Assoc. of Stratigr. Palynol., Dallas, Tex.
- Noormets, R., E. A. Felton, and K. A. W. Crook (2002), Sedimentology of rocky shorelines: 2. Shoreline megaclasts on the north shore of Oahu, Hawaii—Origins and history, *Sediment. Geol.*, *150*, 31–45, doi:10.1016/S0037-0738(01)00266-4.
- Rice, J. W., Jr., and K. S. Edgett (1997), Catastrophic flood sediments in Chryse Basin, Mars, and Quincy Basin, Washington: Application of sandar facies model, *J. Geophys. Res.*, *102*(E2), 4185–4200, doi:10.1029/96JE02824.
- Richardson, K., and P. A. Carling (2005), A typology of sculpted forms in bedrock channels, *Geol. Soc. Am. Spec. Pap. SPE392*, Geol. Soc. of Am., Boulder, Colo.
- Rouai, M., and A. Dekayir (2001), Caractérisation fractale de la porosité d'un basalte altéré, *C. R. Acad. Sci. Ser. 2*, *332*(10), 595–600.
- Ryan, R. J., A. M. O'Beirne-Ryan, and M. Zentilli (2005), Rounded cobbles that have not travelled far: Incorporation of coresotones from saprolites in the South Mountain area of southern Nova Scotia, Canada, *Sedimentology*, *52*, 1109–1121, doi:10.1111/j.1365-3091.2005.00730.x.