

Appendix A

[91] Targets observed by the MI were often also observed by Pancam and sometimes by the other IDD instruments as well. Table A1 summarizes MI observations during the first 900 sols of Opportunity's mission and shows the relationship of each MI observation to RAT or Mössbauer (MB) [Klingelhöfer et al., 2003] observations of the same target. The far right column lists the sequence ID and sol of acquisition of Pancam images of the same target. In some cases, the target was modified by the RAT or MB contact but not observed by Pancam after the modification; in these cases the entry is "none."

[92] **Acknowledgments.** We thank the MER rover planners for their outstanding support of Opportunity IDD operations: Brian K. Cooper, Jeff Biesiadecki, Frank Hartman, Scott Maxwell, John Wright, Jeng Yen, Chris Leger, Robert Bonitz, Eric Baumgartner, Khaled Ali, Ashitesy Trebilco, and Mark Maimone. We also thank the NASA Ames MER support team: Matthew Deans, Laurence Edwards, Joel Hagen, Clayton Kunz, David Lees, Randy Sargent, Michael Wagner, and Anne Wright. Software tools developed at Cornell University by the Pancam team continue to be very useful in tracking and managing MI data products; we thank Elaina McCartney for her repeated assistance in operations planning. The MIPL MER team quickly generated many MI mosaics: Doug Alexander, Amy Chen, Oleg Pariser, Bob Deen, Jeff Hall, Mike Cayan, Vadim Klochko, Elmain Martinez, and Charles Thompson. We also appreciate the support received from the MER team at the USGS in Flagstaff: Jeff Anderson, Tammy Becker, Chris Isbell, Brian Lipkowitz, Dave MacKinnon, Jac Shinaman, Deborah Soltesz, Sean Varga, and Robert Wallace. Boris Semenov of JPL's Navigation and Ancillary Information Facility supported MI geometric processing by providing SPICE kernels in a timely fashion. This work was performed for the Jet Propulsion Laboratory, California Institute of Technology, sponsored by the National Aeronautics and Space Administration. The use of trade, product, or firm names in this paper does not imply endorsement by the U.S. Government.

References

- Ahlbrandt, T. S. (1979), Textural parameters of eolian deposits, *U.S. Geol. Surv. Prof. Pap.*, 1052, 21–52.
- Alexander, D. A., et al. (2006), Processing of Mars Exploration Rover imagery for science and operations planning, *J. Geophys. Res.*, 111, E02S02, doi:10.1029/2005JE002462.
- Arvidson, R. E., et al. (2004), Localization and physical property experiments conducted by Opportunity at Meridiani Planum, *Science*, 306, 1730–1733, doi:10.1126/science.1104211.
- Bell, J. F., III, et al. (2003), Mars Exploration Rover Athena Panoramic Camera (Pancam) investigation, *J. Geophys. Res.*, 108(E12), 8063, doi:10.1029/2003JE002070.
- Bell, J. F., III, J. Joseph, J. N. Sohl-Dickstein, H. M. Arneson, M. J. Johnson, M. T. Lemmon, and D. Savransky (2006), In-flight calibration and performance of the Mars Exploration Rover Panoramic Camera (Pancam) instruments, *J. Geophys. Res.*, 111, E02S03, doi:10.1029/2005JE002444.
- Calvin, W. M., et al. (2008), Hematite spherules at Meridiani: Results from MI, Mini-TES, and Pancam, *J. Geophys. Res.*, doi:10.1029/2007JE003048, in press.
- Carlson, J., and J. P. Grotzinger (2001), Submarine fan environment inferred from turbidite thickness distributions, *J. Sediment. Res.*, 48, 1331–1351, doi:10.1046/j.1365-3091.2001.00426.x.
- Choquette, P. W., and L. C. Pray (1970), Geologic nomenclature and classification of porosity in sedimentary carbonates, *Am. Assoc. Pet. Geol. Bull.*, 54, 207–250.
- Clark, B. C., et al. (2005), Chemistry and mineralogy of outcrops at Meridiani Planum, *Earth Planet. Sci. Lett.*, 240, 73–94, doi:10.1016/j.epsl.2005.09.040.
- Connolly, H. C., Jr., et al. (2006), The meteoritical bulletin, no. 90, 2006 September, *Meteorit. Planet. Sci.*, 41(9), 1383–1418.
- Crisp, J. A., M. Adler, J. R. Matijevic, S. W. Squyres, R. E. Arvidson, and D. M. Kass (2003), Mars Exploration Rover mission, *J. Geophys. Res.*, 108(E12), 8061, doi:10.1029/2002JE002038.
- Gorevan, S. P., et al. (2003), Rock Abrasion Tool: Mars Exploration Rover mission, *J. Geophys. Res.*, 108(E12), 8068, doi:10.1029/2003JE002061.
- Greeley, R., and J. D. Iversen (1985), *Wind as a Geological Process*, 333 pp., Cambridge Univ. Press, Cambridge, U. K.
- Grotzinger, J. P., et al. (2005), Stratigraphy and sedimentology of a dry to wet eolian depositional system, Burns formation, Meridiani Planum, Mars, *Earth Planet. Sci. Lett.*, 240, 11–72, doi:10.1016/j.epsl.2005.09.039.
- Grotzinger, J., et al. (2006), Sedimentary textures formed by aqueous processes, Erebus Crater, Meridiani Planum, Mars, *Geology*, 34, 1085–1088, doi:10.1130/G22985A.1.
- Herkenhoff, K. E., et al. (2003), Athena Microscopic Imager investigation, *J. Geophys. Res.*, 108(E12), 8065, doi:10.1029/2003JE002076.
- Herkenhoff, K., et al. (2004a), Mars Exploration Rover Project Microscopic Imager calibration report, *Rev. B. D-19830, MER 420–6-704*, Jet Propul. Lab., Pasadena, Calif.
- Herkenhoff, K. E., et al. (2004b), Evidence from Opportunity's Microscopic Imager for water on Meridiani Planum, *Science*, 306, 1727–1730, doi:10.1126/science.1105286.
- Herkenhoff, K. E., et al. (2006), Overview of the Microscopic Imager investigation during Spirit's first 450 sols in Gusev Crater, *J. Geophys. Res.*, 111, E02S04, doi:10.1029/2005JE002574.
- Jolliff, B. L., W. H. Farrand, J. R. Johnson, C. Schröder, C. M. Weitz, and the Athena Science Team (2006), Origin of rocks and cobbles on the Meridiani Plains as seen by Opportunity, *Lunar Planet. Sci.*, XXXVII, Abstract 2401.
- Klingelhöfer, G., et al. (2003), The Athena MIMOS II Mössbauer spectrometer investigation, *J. Geophys. Res.*, 108(E12), 8067, doi:10.1029/2003JE002138.
- Klingelhöfer, G., et al. (2004), Jarosite and hematite at Meridiani Planum from Opportunity's Mössbauer spectrometer, *Science*, 306, 1740–1746, doi:10.1126/science.1104653.
- Knauth, L. P., D. M. Burt, and K. H. Wohletz (2005), Impact origin of sediments at the Opportunity landing site on Mars, *Nature*, 438, 1123–1128, doi:10.1038/nature04383.
- Knoll, A. H., et al. (2008), Veneers, rinds, and fracture fills: Relatively late alteration of sedimentary rocks at Meridiani Planum, Mars, *J. Geophys. Res.*, 113, E06S16, doi:10.1029/2007JE002949.
- Lewis, K., A. Aharonson, J. P. Grotzinger, and S. W. Squyres (2008), Structure and stratigraphy of Home Plate from the Spirit Mars Exploration Rover, *J. Geophys. Res.*, doi:10.1029/2007JE003025, in press.
- Maki, J. N., et al. (2003), The Mars Exploration Rover engineering cameras, *J. Geophys. Res.*, 108(E12), 8071, doi:10.1029/2003JE002077.
- McCullom, T. M., and B. M. Hynek (2005), A volcanic environment for bedrock diagenesis at Meridiani Planum on Mars, *Nature*, 438, 1129–1131, doi:10.1038/nature04390.
- McLennan, S. M., et al. (2005), Provenance and diagenesis of the evaporate-bearing Burns formation, Meridiani Planum, Mars, *Earth Planet. Sci. Lett.*, 240, 95–121, doi:10.1016/j.epsl.2005.09.041.
- Metz, J. M., J. P. Grotzinger, D. M. Rubin, K. Lewis, S. W. Squyres, and J. F. Bell III (2008), Sulfate-rich eolian and wet interdune deposits, Erebus Crater, Meridiani Planum, Mars, *J. Sediment. Res.*, in press.
- Morris, R. V., et al. (2006), Mössbauer mineralogy of rock, soil, and dust at Meridiani Planum, Mars: Opportunity's journey across sulfate-rich outcrop, basaltic sand and dust, and hematite lag deposits, *J. Geophys. Res.*, 111, E12S15, doi:10.1029/2006JE002791.
- Perl, S. M., S. M. McLennan, J. P. Grotzinger, K. E. Herkenhoff, and the Athena Science Team (2007), Sedimentological constraints on an infiltrating paleowater table in the Burns formation, Meridiani Planum, Mars, *Seventh International Conference on Mars*, Abstract 3298, Jet Propul. Lab., Pasadena, Calif.
- Powers, M. (1953), A new roundness scale for sedimentary particles, *J. Sediment. Res.*, 23, 117–119.
- Reid, R. J., et al. (1999), Imager for Mars Pathfinder (IMP) image calibration, *J. Geophys. Res.*, 104, 8907–8926, doi:10.1029/1998JE900011.
- Rothman, D. H., and J. P. Grotzinger (1995), Scaling properties of gravity-driven sediments, *Nonlinear Processes Geophys.*, 2, 178–185.
- Schröder, C., et al. (2008), Meteorites on Mars observed with the Mars Exploration Rovers, *J. Geophys. Res.*, 113, E06S22, doi:10.1029/2007JE002990.
- Soderblom, J. M., J. F. Bell III, J. R. Johnson, J. Joseph, and M. J. Wolff (2008), Mars Exploration Rover Navigation Camera in-flight calibration, *J. Geophys. Res.*, 113, E06S19, doi:10.1029/2007JE003003.
- Soderblom, L. A., et al. (2004), Soils of Eagle Crater and Meridiani Planum at the Opportunity Rover landing site, *Science*, 306, 1723–1726, doi:10.1126/science.1105127.
- Southard, J. B., and L. A. Boguchwal (1990), Bed configuration in steady unidirectional water flows: Part 2. Synthesis of flume data, *J. Sediment. Res.*, 60, 658–679.
- Sparks, R. S. J. (1976), Grain-size variations in ignimbrites and implications for transport of pyroclastic flows, *Sedimentology*, 23, 147–188, doi:10.1111/j.1365-3091.1976.tb00045.x.
- Squyres, S. W., and A. H. Knoll (2005), Sedimentary rocks at Meridiani Planum: Origin, diagenesis, and implications for life on Mars, *Earth Planet. Sci. Lett.*, 240, 1–10, doi:10.1016/j.epsl.2005.09.038.