

- Mangold, N., Mangeney, A., Bouchut, F., 2008a. Levee-channel deposits in dry or wet debris flows: a tool to understand gullies formation, in Workshop on Martian Gullies: Theories and Tests, Abstract #8006, League City, TX.
- Mangold, N., Mangeney, A., Bouchut, F., Ionescu, I., 2008b. The key is in the bend: how sinuous gullies improve the understanding of their flow properties, in American Geophysical Union Fall Meeting, Abstract #P34A-02, San Francisco, CA.
- Marchant, D.R., Head, J.W., 2007. Antarctic Dry Valleys: microclimate zonation, variable geomorphic processes, and implications for assessing climate change on Mars. *Icarus* 192 (1), 187–222.
- McEwen, A.S., Eliason, E.M., Bergstrom, J.W., Bridges, N.T., Hansen, C.J., Delamere, W.A., Grant, J.A., Gulick, V.C., Herkenhoff, K.E., Keszthelyi, L., Kirk, R.L., Mellon, M.T., Squyres, S.W., Thomas, N., Weitz, C.M., 2007. A closer look at water-related geologic activity on Mars. *Science* 317 (5845), 1706–1709.
- McEwen, A.S., Banks, M.E., Baugh, N., Becker, K., Boyd, A., Bergstrom, J.W., Beyer, R.A., Bortolini, E., Bridges, N.T., Byrne, S., Castalla, B., Chuang, F.C., Crumpler, L.S., Daubar, I., Davatzes, A.K., Deardorff, D.G., DeJong, E., Delamere, W.A., Dobrea, E.Z.N., Dundas, C.M., Eliason, E.M., Espinoza, Y., Fennema, A., Fishbaugh, K.E., Forrester, T., Geissler, P.E., Grant, J.A., Griffes, J.L., Grotzinger, J.P., Gulick, V.C., Hansen, C.J., Herkenhoff, K.E., Heyd, R., Jaeger, W.L., Jones, D., Kanefsky, B., Keszthelyi, L., King, R., Kirk, R.L., Kolb, K.J., Lasco, J., Lefort, A., Leis, R., Lewis, K.W., Martinez-Alonso, S.E., Mattson, S., McArthur, G., Mellon, M.T., Metz, J.M., Milazzo, M.P., Milliken, R.E., Motazedian, T., Okubo, C.H., Ortiz, A., Phillippoff, A.J., Plassmann, J., Polit, A., Russell, P.S., Schaller, C., Searis, M.L., Spriggs, T., Squyres, S.W., Tarr, S., Thomas, N., Thomson, B.J., Tornabene, L.L., Houten, C.V., Verba, C., Weitz, C.M., Wray, J.J., 2010. The High Resolution Imaging Science Experiment (HiRISE) during MRO's primary science phase. *Icarus*. *Nature* 205 (1), 2–37.
- Milliken, R.E., Mustard, J.F., Goldsby, D.L., 2003. Viscous flow features on the surface of Mars: observations from high-resolution Mars Orbiter Camera (MOC) images. *J. Geophys. Res.* 108 (E6). doi:10.1029/2002JE002005.
- Miyamoto, H., Dohm, J.M., Baker, V.R., Beyer, R.A., Bourke, M., 2004. Dynamics of unusual debris flows on Martian sand dunes. *Geophys. Res. Lett.* 31. doi:10.1029/2004GL020313.
- Morgan, G.A., Head, J.W., Marchant, D.R., Dickson, J.L., Levy, J.S., 2007a. Gully formation on Mars: Testing the snowpack hypothesis from analysis of analogs in the Antarctic Dry Valleys, in Seventh International Conference on Mars, Abstract #1353, Pasadena, CA.
- Morgan, G.A., Head, J.W., Marchant, D.R., Dickson, J.L., Levy, J.S., 2007b. Gully formation on Mars: Testing the snowpack hypothesis from analysis of analogs in the Antarctic Dry Valleys, in LPSC 38, Abstract #1656, League City, TX.
- Morgan, G.A., Head, J.W., Marchant, D.R., Dickson, J.L., Levy, J.S., 2008. Gully formation and evolution in the Antarctic Dry Valleys: Implications for Mars, in Workshop on Martian Gullies, Abstract #1301, Houston, TX.
- Mustard, J.F., Cooper, C.D., Rifkin, M.K., 2001. Evidence for recent climate change on Mars from the identification of youthful near-surface ground ice. *Nature* 412 (6845), 411–414.
- Neukum, G., Jaumann, R., the HRSC Co-Investigator and ExperimentTeam, 2004. HRSC: the high resolution stereo camera of Mars express. European Space Agency Special Publication, ESA SP-1240, pp. 17–35.
- Oberst, J., Scholten, F., Gwinner, K., Giese, B., Waehlich, M., Roatsch, T., Neukum, G., 2005. The topographic mapping performance of the HRSC (High Resolution Stereo Camera) on Mars Express, in American Geophysical Union Fall Meeting, Abstract #G53C-03, San Francisco, CA.
- Parsons, R.A., Nimmo, F., Kreslavsky, M., 2008. Fluvial discharge rates of martian gullies: slope measurements from stereo HiRISE images and numerical modeling of sediment transport, in American Geophysical Union Fall Meeting, Abstract #P41A-1350, San Francisco, CA.
- Pelletier, J.D., Kolb, K.J., McEwen, A.S., Kirk, R.L., 2008. Recent bright gully deposits on Mars: wet or dry flow? *Geology* 36 (3), 211–214.
- Richter, L., Schmitz, N., Weiss, S., the MER/Athena Team, 2006. Inference of strength of soil deposits along MER rover traverses, in European Planetary Science Conference, p. 523, Berlin, Germany.
- Riess, D., van Gasselt, S., Neukum, G., Jaumann, R., 2004. Absolute dune ages and implications for the time of formation of gullies in Nirgal Valles, Mars. *J. Geophys. Res.* 109. doi:10.1029/2004JE002251.
- Schon, S.C., Head, J.W., Fassett, C.I., 2009. Unique chronostratigraphic marker in depositional fan stratigraphy on Mars: evidence for ~1.25 Ma old gully activity and surficial meltwater origin. *Geology* 37. doi:10.1130/G25398A.25391.
- Sharp, R.P., 1973. Mars: fretted and chaotic terrains. *J. Geophys. Res.* 78, 4073–4083.
- Shinbrot, T., Duong, N.-H., Kwan, L., Alvarez, M.M., 2004. Dry granular flows can generate surface features resembling those seen in martian gullies. *Proc. Natl. Acad. Sci.* 101 (23), 8542–8546.
- Treiman, A.H., 2003. Geologic settings of martian gullies: implications for their origins. *J. Geophys. Res.* 108. doi:10.1029/2002JE001900.
- Williams, K.E., Toon, O.B., Heldmann, J.L., Mellon, M.T., 2009. Ancient melting of mid-latitude snowpacks on Mars as a water source for gullies. *Icarus* 199. doi:10.1016/j.icarus.2008.1012.1013.