

Supporting information part (II) of (II)

Synthesis of 2',4'-Propylene-Bridged (carba-ENA) Thymidine and its analogues - The Engineering of Electrostatic and Steric Effects at The Bottom of the Minor Groove for Nuclease and Thermodynamic Stabilities and Elicitation of RNase H

Yi Liu, Jianfeng Xu, Mansoureh Karimiahmadabadi, and Jyoti Chattopadhyaya*

Bioorganic Chemistry Program, Department of Cell and Molecular Biology, Box 581, Biomedical Center,
Uppsala University, SE-751 23 Uppsala, Sweden

Jyoti@boc.uu.se

Figure SII. 1. 1D nOe spectrum of compound 14 (H1' irradiated).....	3
Figure SII. 2. 1D nOe spectrum of compound 20a (H1' irradiated)	4
Figure SII. 3. 1D nOe spectrum of compound 21b (H8' irradiated).....	5
Figure SII. 4. 1D nOe spectrum of compound 26 (H1' irradiated).....	6
Figure SII. 5. 1D nOe spectrum of compound 28 (H-1' _{8' S} irradiated).....	7
Figure SII. 6. 1D nOe spectrum of compound 28 (H-1' _{8' R} irradiated)	8
Figure SII. 7. Homodecoupling spectrum of compound 14.....	9
Figure SII. 8. Homodecoupling spectrum of compound 16b.....	10
Figure SII. 9. Homodecoupling spectrum of compound 16b (H6' decoupled).....	11
Figure SII. 10. Homodecoupling spectrum of compound 21a.....	12
Figure SII. 11. Double Homodecoupling spectrum of compound 21a (H8' and H6' ' decoupled)	13
Figure SII. 12. Homodecoupling spectrum of compound 21b.....	14
Figure SII. 13. Double Homodecoupling spectrum of compound 21b (H8' and H6' ' decoupled).....	15