

REFERENCES

- Albin RL, Young AB, and Penney JB.** The functional anatomy of disorders of the basal ganglia. *Trends in Neurosciences* 18: 63-63, 1995.
- Alexander GE, DeLong MR, and Strick PL.** Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Ann Rev Neurosci* 9: 357-381, 1986.
- Amador S, C., Hood A, J., Schiess MC, Izor R, and Sereno A, B.** Dissociating cognitive deficits involved in voluntary eye movement dysfunctions in Parkinson's disease patients. *Neuropsychologia* 44: 1475-1482, 2006.
- Anderson ME, Postupna N, and Ruffo M.** Effects of High-Frequency Stimulation in the Internal Globus Pallidus on the Activity of Thalamic Neurons in the Awake Monkey. *J Neurophysiol* 89: 1150-1160, 2003.
- Ashby P, Kim YJ, Kumar R, Lang AE, and Lozano AM.** Neurophysiological effects of stimulation through electrodes in the human subthalamic nucleus. *Brain* 122: 1919-1931, 1999.
- Basso MA and Evinger C.** An explanation for reflex blink hyperexcitability in Parkinson's disease. II. Nucleus raphe magnus. *J Neurosci* 16: 7318-7330, 1996.
- Basso MA, Pokorny JJ, and Liu P.** Activity of monkey substantia nigra pars reticulata neurons during smooth pursuit eye movements. *European Journal of Neuroscience* 22: 448-464, 2005.
- Basso MA, Powers AS, and Evinger C.** An explanation for reflex blink hyperexcitability in Parkinson's disease. I. Superior Colliculus. *J Neurosci* 16: 7308-7317, 1996.
- Basso MA, R.E. S, and Evinger C.** Midbrain 6-hydroxydopamine lesions modulate blink reflex excitability. *Exp Brain Res* 94: 88-96, 1993.
- Basso MA and Wurtz RH.** Neuronal activity in substantia nigra pars reticulata during target selection. *J. Neuroscience* 22: 1883-1894, 2002.
- Bayer HM, Handel A, and Glimcher PW.** Eye position and memory saccade related responses in substantia nigra pars reticulata. *Experimental Brain Research* 154: 428 - 441, 2002.
- Beckstead RM.** Long collateral branches of substantia nigra pars reticulata axons to thalamus, superior colliculus and reticular formation in monkey and cat. Multiple retrograde neuronal labeling with fluorescent dyes. *Neurosci* 10: 767-779, 1983.
- Beckstead RM, Edwards SB, and Frankfurter A.** A comparison of the intranigral distribution of nigrotectal neurons labeled with horseradish peroxidase in the monkey, cat, and rat. *J Neurosci* 1: 121-125, 1981.
- Behan M and Kime NM.** Intrinsic circuitry in the deep layers of the cat superior colliculus. *Visual Neuroscience* 13: 1031-1042, 1996.
- Behan M, Steinhacker K, Jeffrey-Borger S, and Meredith M.** Chemoarchitecture of GABAergic neurons in the ferret superior colliculus. *Journal of Comparative Neurology* 452: 334-359, 2002.
- Benabid AL.** Deep brain stimulation for Parkinson's disease. *Current Opinion in Neurobiology* 13: 696-706, 2003.