

Supplemental references

1. Park, S. et al. Hexameric assembly of the proteasomal ATPases is templated through their C termini. *Nature* 459, 866-70 (2009).
2. Groll, M. et al. Structure of 20S proteasome from yeast at 2.4 Å resolution. *Nature* 386, 463-71 (1997).
3. Zhang, F. et al. Structural insights into the regulatory particle of the proteasome from *Methanocaldococcus jannaschii*. *Molecular cell* 34, 473-84 (2009).
4. Djuranovic, S. et al. Structure and activity of the N-terminal substrate recognition domains in proteasomal ATPases. *Molecular cell* 34, 580-90 (2009).
5. Saeki, Y., Toh, E.A., Kudo, T., Kawamura, H. & Tanaka, K. Multiple proteasome-interacting proteins assist the assembly of the yeast 19S regulatory particle. *Cell* 137, 900-13 (2009).
6. Kaneko, T. et al. Assembly pathway of the Mammalian proteasome base subcomplex is mediated by multiple specific chaperones. *Cell* 137, 914-25 (2009).
7. Funakoshi, M., Tomko, R.J., Jr., Kobayashi, H. & Hochstrasser, M. Multiple assembly chaperones govern biogenesis of the proteasome regulatory particle base. *Cell* 137, 887-99 (2009).
8. Leggett, D.S. et al. Multiple associated proteins regulate proteasome structure and function. *Molecular cell* 10, 495-507 (2002).
9. Finley, D., Ozkaynak, E. & Varshavsky, A. The yeast polyubiquitin gene is essential for resistance to high temperatures, starvation, and other stresses. *Cell* 48, 1035-46 (1987).