

## REFERENCES

1. Hurley, J. H. (2008). ESCRT complexes and the biogenesis of multivesicular bodies. *Curr Opin Cell Biol* **20**, 4-11.
2. Saksena, S., Sun, J., Chu, T. & Emr, S. D. (2007). ESCRTing proteins in the endocytic pathway. *Trends Biochem Sci* **32**, 561-73.
3. Williams, R. L. & Urbe, S. (2007). The emerging shape of the ESCRT machinery. *Nat Rev Mol Cell Biol* **8**, 355-68.
4. Welsch, S., Habermann, A., Jager, S., Muller, B., Krijnse-Locker, J. & Krausslich, H. G. (2006). Ultrastructural analysis of ESCRT proteins suggests a role for endosome-associated tubular-vesicular membranes in ESCRT function. *Traffic* **7**, 1551-66.
5. Hanson, P. I., Roth, R., Lin, Y. & Heuser, J. E. (2008). Plasma membrane deformation by circular arrays of ESCRT-III protein filaments. *J Cell Biol* **180**, 389-402.
6. Babst, M., Sato, T. K., Banta, L. M. & Emr, S. D. (1997). Endosomal transport function in yeast requires a novel AAA-type ATPase, Vps4p. *Embo J* **16**, 1820-31.
7. Carlton, J. G. & Martin-Serrano, J. (2007). Parallels between cytokinesis and retroviral budding: a role for the ESCRT machinery. *Science* **316**, 1908-12.
8. Morita, E., Sandrin, V., Chung, H. Y., Morham, S. G., Gygi, S. P., Rodesch, C. K. & Sundquist, W. I. (2007). Human ESCRT and ALIX proteins interact with proteins of the midbody and function in cytokinesis. *Embo J* **26**, 4215-27.
9. Finken-Eigen, M., Rohricht, R. A. & Kohrer, K. (1997). The VPS4 gene is involved in protein transport out of a yeast pre-vacuolar endosome-like compartment. *Curr Genet* **31**, 469-80.
10. Fujita, H., Yamanaka, M., Imamura, K., Tanaka, Y., Nara, A., Yoshimori, T., Yokota, S. & Himeno, M. (2003). A dominant negative form of the AAA ATPase SKD1/VPS4 impairs membrane trafficking out of endosomal/lysosomal compartments: class E vps phenotype in mammalian cells. *J Cell Sci* **116**, 401-14.
11. Yoshimori, T., Yamagata, F., Yamamoto, A., Mizushima, N., Kabeya, Y., Nara, A., Miwako, I., Ohashi, M., Ohsumi, M. & Ohsumi, Y. (2000). The mouse SKD1, a homologue of yeast Vps4p, is required for normal endosomal trafficking and morphology in mammalian cells. *Mol Biol Cell* **11**, 747-63.
12. Bishop, N. & Woodman, P. (2000). ATPase-defective mammalian VPS4 localizes to aberrant endosomes and impairs cholesterol trafficking. *Mol Biol Cell* **11**, 227-39.
13. Fujii, K., Hurley, J. H. & Freed, E. O. (2007). Beyond Tsg101: the role of Alix in 'ESCRTing' HIV-1. *Nat Rev Microbiol* **5**, 912-6.
14. Welsch, S., Muller, B. & Krausslich, H. G. (2007). More than one door - Budding of enveloped viruses through cellular membranes. *FEBS Lett* **581**, 2089-97.
15. Bieniasz, P. D. (2006). Late budding domains and host proteins in enveloped virus release. *Virology* **344**, 55-63.