

# References

1. Girou E, Loyeau S, Legrand P, Oppein F, Brun-Buisson C: **Efficacy of handrubbing with alcohol based solution versus standard handwashing with antiseptic soap: randomised clinical trial.** *Bmj* 2002, **325(7360):**362.
2. Austin DJ, Bonten MJ, Weinstein RA, Slaughter S, Anderson RM: **Vancomycin-resistant enterococci in intensive-care hospital settings: transmission dynamics, persistence, and the impact of infection control programs.** *Proc Natl Acad Sci USA* 1999, **96(12):**6908-6913.
3. McBryde ES, Pettitt AN, McElwain DL: **A stochastic mathematical model of methicillin resistant *Staphylococcus aureus* transmission in an intensive care unit: predicting the impact of interventions.** *J Theor Biol* 2007, **245(3):**470-481.
4. J.M. Boyce and D. Pittet, "Guidelines for Hand Hygiene in Health-Care Settings: Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force," *Infection Control and Hospital Epidemiology* 23 (2002), pp. S3-S41.
5. D. Pittet, B. Allegranzi, and J.M. Boyce, "The World Health Organization Guidelines on Hand Hygiene in Health Care and Their Consensus Recommendations," *Infection Control and Hospital Epidemiology* 30, World Health Organization World Alliance for Patient Safety First Global Patient Safety Challenge Core Group of Experts (2009), pp. 611-622.
6. J.P. Haas and E.L. Larson, "Measurement of Compliance with Hand Hygiene," *Journal of Hospital Infection* 66 (2007), pp. 6-14.