

3.1	Degree distributions for the <i>sparse</i> ₅₀ HCW contact network and corresponding graphs generated by the Erdős-Rényi (ER), Chung-Lu (CL), and Barabási-Albert (BA) models. <i>x</i> -axis denotes the degree. <i>y</i> -axis denotes the fraction of nodes. Plot is truncated; maximum values can be found in Table 3.1.	52
3.2	Degree distributions for the <i>moderate</i> ₅₀ HCW contact network and corresponding graphs generated by the Erdős-Rényi (ER), Chung-Lu (CL), and Barabási-Albert (BA) models. <i>x</i> -axis denotes the degree. <i>y</i> -axis denotes the fraction of nodes. Plot is truncated; maximum values can be found in Table 3.2.	53
3.3	Comparison of graphs generated by the Erdős-Rényi (ER), Barabási-Albert (BA), Chung-Lu (CL), Configuration (CONF), and Configuration with Assortativity (CA) models with the <i>moderate</i> ₅₀ (HCW) graph they model. Graph shows the mean number of people infected on a given day of the SIR simulation with transmissibility ($\rho = .0302$). <i>x</i> -axis represents the timestep for the simulation (days). <i>y</i> -axis gives the number of people in the “infected” state during that timestep.	57
3.4	Comparison of graphs generated by the Erdős-Rényi (ER), Barabási-Albert (BA), Chung-Lu (CL), Configuration (CONF), and Configuration with Assortativity (CA) models with the <i>moderate</i> ₅₀ (HCW) graph they model. Graph shows the median number of people infected on a given day of the SIR simulation with transmissibility ($\rho = .0302$). <i>x</i> -axis represents the timestep for the simulation (days). <i>y</i> -axis gives the number of people in the “infected” state during that timestep.	58
3.5	Comparison of graphs generated by the Erdős-Rényi (ER), Barabási-Albert (BA), Chung-Lu (CL), Configuration (CONF), and Configuration with Assortativity (CA) models with the <i>moderate</i> ₅₀ (HCW) graph they model. Graph shows the mean number of people infected on a given day of the SIR simulation with transmissibility $\rho = .0122$. <i>x</i> -axis represents the timestep for the simulation (days). <i>y</i> -axis gives the number of people in the “infected” state during that timestep.	61
3.6	Comparison of the HCW contact network and CONF graphs for transmissibility values in $\{.00763, 0.00794, 0.00824, 0.00854\}$	64
3.7	Comparison of the HCW contact network and CONF graphs for transmissibility values in $\{.00763, 0.00794, 0.00824, 0.00854\}$	66
3.8	Clustering coefficient distribution for the <i>sparse</i> ₅₀ , <i>moderate</i> ₅₀ , and <i>dense</i> ₅₀ HCW contact networks. Vertices with degree less than 2 are ignored. . .	69