

- Modell., 221, 2458–2474, doi:10.1016/j.ecolmodel.2010.07.008, 2010.
- Betts, A. K. and Ball, J. H.: Albedo over the boreal forest, *J. Geophys. Res.*, 102, 28901–28909, 1997.
- Betts, R. A.: Offset of the potential carbon sink from boreal forestation by decreases in surface albedo., *Nature*, 408, 187–90, doi:10.1038/35041545, 2000.
- Bonan, G. B., Lawrence, P. J., Oleson, K. W., Levis, S., Jung, M., Reichstein, M., Lawrence, D. M., and Swenson, S. C.: Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data, *J. Geophys. Res.*, 116, 1–22, doi:10.1029/2010JG001593, 2011.
- Bréda, N. J. J.: Ground-based measurements of leaf area index: a review of methods, instruments and current controversies, *J. Experiment. Botan.*, 54, 2403–17, doi:10.1093/jxb/erg263, 2003.
- Bright, R. M., Cherubini, F., and Strømman, A. H.: Climate impacts of bioenergy: Inclusion of carbon cycle and albedo dynamics in life cycle impact assessment, *Environ. Impact Assess. Rev.*, 37, 2–11, doi:10.1016/j.eiar.2012.01.002, 2012.
- Brus, D. J., Hengeveld, G. M., Walvoort, D. J. J., Goedhart, P. W., Heidema, A. H., Nabuurs, G. J., and Gunia, K.: Statistical mapping of tree species over Europe, *Europ. J. Forest Res.*, 131, 145–157, doi:10.1007/s10342-011-0513-5, 2011.
- Cescatti, A., Marcolla, B., Santhana Vannan, S. K., Pan, J. Y., Román, M. O., Yang, X., Ciaïis, P., Cook, R. B., Law, B. E., Matteucci, G., Migliavacca, M., Moors, E., Richardson, A. D., Seufert, G. and Schaaf, C. B.: Intercomparison of MODIS albedo retrievals and in situ measurements across the global FLUXNET network, *Remote Sens. Environ.*, 121, 323–334, doi:10.1016/j.rse.2012.02.019, 2012.
- Chen, J. M.: Optically-based methods for measuring seasonal variation of leaf area index in boreal conifer stands, *Agr. Forest Meteorol.*, 80, 135–163, doi:10.1016/0168-1923(95)02291-0, 1996.
- Cherubini, F., Bright, R. M., and Strømman, A. H.: Site-specific global warming potentials of biogenic CO₂ for bioenergy: contributions from carbon fluxes and albedo dynamics, *Environ. Res. Lett.*, 7, 045902, doi:10.1088/1748-9326/7/4/045902, 2012.
- Condés, S. and Sterba, H.: Derivation of compatible crown width equations for some important tree species of Spain, *Forest Ecol. Manage.*, 217, 203–218, doi:10.1016/j.foreco.2005.06.002, 2005.
- Davidson, A. and Wang, S.: The effects of sampling resolution on the surface albedos of dominant land cover types in the North American boreal region, *Remote Sens. Environ.*, 93, 211–224, doi:10.1016/j.rse.2004.07.005, 2004.
- Dickinson, R. E., Henderson-Sellers, A., and Kennedy, P. J.: Biosphere-Atmosphere Transfer Scheme (BATS) Version 1e as Coupled to the NCAR Community Climate Model, NCAR Technical Note, NCAR/TN-38(August), 1993.
- Dore, S., Montes-Helu, M., Hart, S. C., Hungate, B. a., Koch, G. W., Moon, J. B., Finkral, A. J. and Kolb, T. E.: Recovery of ponderosa pine ecosystem carbon and water fluxes from thinning and stand-replacing fire, *Global Change Biol.*, 18, 3171–3185, doi:10.1111/j.1365-2486.2012.02775.x, 2012.
- Fontes, L., Bontemps, J., and Bugmann, H.: Models for supporting forest management in a changing environment, *Forest Systems*, 19, 8–29, 2010.
- Foré, S. A., Vankat, J. L., and Schaefer, R. L.: Temporal variation in the woody understory of an old-growth *Fagus-Acer* forest and implications for overstory recruitment, *J. Veget. Sci.*, 8, 607–614, doi:10.2307/3237364, 1997.
- Gao, F.: MODIS bidirectional reflectance distribution function and albedo Climate Modeling Grid products and the variability of albedo for major global vegetation types, *J. Geophys. Res.*, 110, D01104, doi:10.1029/2004JD005190, 2005.
- Hansen, J., Sato, M., and Ruedy, R.: Radiative forcing and climate response, *J. Geophys. Res.-Atmos.*, 102, 6831–6864, doi:10.1029/96JD03436, 1997.
- Haverd, V., Lovell, J. L., Cuntz, M., Jupp, D. L. B., Newnham, G. J., and Sea, W.: The Canopy Semi-analytic Pgap And Radiative Transfer (CanSPART) model: Formulation and application, *Agric. For. Meteorol.*, 160, 14–35, doi:10.1016/j.agrformet.2012.01.018, 2012.
- Hollinger, D. Y., Ollinger, S. V., Richardson, a. D., Meyers, T. P., Dail, D. B., Martin, M. E., Scott, N. A., Arkebauer, T. J., Baldocchi, D. D., Clark, K. L., Curtis, P. S., Davis, K. J., Desai, A. R., Dragoni, D., Goulden, M. L., Gu, L., Katul, G. G., Pallardy, S. G., Paw U, K. T., Schmid, H. P., Stoy, P. C., Suyker, A. E., and Verma, S. B.: Albedo estimates for land surface models and support for a new paradigm based on foliage nitrogen concentration, *Glob. Change Biol.*, 16, 696–710, doi:10.1111/j.1365-2486.2009.02028.x, 2010.
- Houspanossian, J., Nosetto, M., and Jobbágy, E. G.: Radiation budget changes with dry forest clearing in temperate Argentina, *Glob. Change Biol.*, 19, 1211–1222, doi:10.1111/gcb.12121, 2013.
- Jansen, J. J., Sevenster, J., and Faber, J.: Opbrensttabellen voor belangrijke boomsoorten in Nederland, IBN-rapport 221. Hinkelord Report 17. DLO-Instituut voor Bos- en Natuuronderzoek; Landbouwniversiteit, Wageningen. 202 p, IBN, Instituut voor Bos- en Natuuronderzoek, Institute for Forestry and Nature Research., 1996.
- Kirschbaum, M. U. F., Whitehead, D., Dean, S. M., Beets, P. N., Shepherd, J. D., and Ausseil, A.-G. E.: Implications of albedo changes following afforestation on the benefits of forests as carbon sinks, *Biogeosciences*, 8, 3687–3696, doi:10.5194/bg-8-3687-2011, 2011.
- Kobayashi, H., Ryu, Y., Baldocchi, D. D., Welles, J. M., and Norman, J. M.: On the correct estimation of gap fraction: How to remove scattered radiation in gap fraction measurements?, *Agric. For. Meteorol.*, 174–175, 170–183, doi:10.1016/j.agrformet.2013.02.013, 2013.
- Kramer, K., Buiteveld, J., Forstreuter, M., Geburek, T., Leonardi, S., Menozzi, P., Povillon, F., Schelhaas, M.-J., Teissier du Cros, E., Vendramin, G. G., and Van der Werf, D. C.: Bridging the gap between ecophysiological and genetic knowledge to assess the adaptive potential of European beech, *Ecol. Modell.*, 216, 333–353, doi:10.1016/j.ecolmodel.2008.05.004, 2008.
- Krinner, G.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochem. Cycles*, 19, doi:10.1029/2003GB002199, 2005.
- Kuusik, A. and Nilson, T.: A directional multispectral forest reflectance model, *Remote Sens. Environ.*, 252, 244–252, 2000.
- Kuusik, A., Kuusik, J., and Lang, M.: A dataset for the validation of reflectance models, *Remote Sens. Environ.*, 113, 889–892, doi:10.1016/j.rse.2009.01.005, 2009.