

- Criscuolo A, Gribaldo S. 2010. BMGE (Block Mapping and Gathering with Entropy): a new software for selection of phylogenetic informative regions from multiple sequence alignments. *BMC Evol Biol.* 10:210.
- de la Torre JR, Walker CB, Ingalls AE, Konneke M, Stahl DA. 2008. Cultivation of a thermophilic ammonia oxidizing archaeon synthesizing crenarchaeol. *Environ Microbiol.* 10:810–818.
- DeLong EF. 1992. Archaea in coastal marine environments. *Proc Natl Acad Sci U S A.* 89:5685–5689.
- DeLong EF, et al. 2006. Community genomics among stratified microbial assemblages in the ocean's interior. *Science* 311:496–503.
- Eisen JA. 1998. Phylogenomics: improving functional predictions for uncharacterized genes by evolutionary analysis. *Genome Res.* 8: 163–167.
- Frigaard NU, Martinez A, Mincer TJ, DeLong EF. 2006. Proteorhodopsin lateral gene transfer between marine planktonic Bacteria and Archaea. *Nature* 439:847–850.
- Fuhrman JA, Davis AA. 1997. Widespread Archaea and novel Bacteria from the deep sea as shown by 16S rRNA gene sequences. *Mar Ecol Prog Ser.* 150:275–285.
- Fuhrman JA, McCallum K, Davis AA. 1992. Novel major archaeobacterial group from marine plankton. *Nature* 356:148–149.
- García-Vallve S, Palau J, Romeu A. 1999. Horizontal gene transfer in glycosyl hydrolases inferred from codon usage in *Escherichia coli* and *Bacillus subtilis*. *Mol Biol Evol.* 16:1125–1134.
- Ghai R, et al. 2010. Metagenome of the Mediterranean deep chlorophyll maximum studied by direct and fosmid library 454 pyrosequencing. *ISME J.* 4:1154–1166.
- Gogarten JP, Doolittle WF, Lawrence JG. 2002. Prokaryotic evolution in light of gene transfer. *Mol Biol Evol.* 19:2226–2238.
- Gophna U, Ofra Y. 2011. Lateral acquisition of genes is affected by the friendliness of their products. *Proc Natl Acad Sci U S A.* 108:343–348.
- Heyer LJ, Kruglyak S, Yooshep S. 1999. Exploring expression data: identification and analysis of coexpressed genes. *Genome Res.* 9: 1106–1115.
- Huang Y, Gilna P, Li W. 2009. Identification of ribosomal RNA genes in metagenomic fragments. *Bioinformatics* 25:1338–1340.
- Hyatt D, et al. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119.
- Isambert H, Stein RR. 2009. On the need for widespread horizontal gene transfers under genome size constraint. *Biol Direct.* 4:28.
- Iverson V, et al. 2012. Untangling genomes from metagenomes: revealing an uncultured class of marine Euryarchaeota. *Science* 335: 587–590.
- Kanhere A, Vingron M. 2009. Horizontal gene transfers in prokaryotes show differential preferences for metabolic and translational genes. *BMC Evol Biol.* 9:9.
- Karner MB, DeLong EF, Karl DM. 2001. Archaeal dominance in the mesopelagic zone of the Pacific Ocean. *Nature* 409:507–510.
- Katoh K, Kuma K, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res.* 33: 511–518.
- Konneke M, et al. 2005. Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* 437:543–546.
- Koski LB, Golding GB. 2001. The closest BLAST hit is often not the nearest neighbor. *J Mol Evol.* 52:540–542.
- Leininger S, et al. 2006. Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* 442:806–809.
- López-García P, Brochier C, Moreira D, Rodríguez-Valera F. 2004. Comparative analysis of a genome fragment of an uncultivated mesopelagic crenarchaeote reveals multiple horizontal gene transfers. *Environ Microbiol.* 6:19–34.
- López-García P, Moreira D, López-López A, Rodríguez-Valera F. 2001. A novel haloarchaeal-related lineage is widely distributed in deep oceanic regions. *Environ Microbiol.* 3:72–78.
- Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25: 955–964.
- Marchler-Bauer A, et al. 2005. CDD: a conserved domain database for protein classification. *Nucleic Acids Res.* 33:D192–D196.
- Martin-Cuadrado AB, et al. 2007. Metagenomics of the deep Mediterranean, a warm bathypelagic habitat. *PLoS One* 2:e914.
- Martin-Cuadrado AB, et al. 2008. Hindsight in the relative abundance, metabolic potential and genome dynamics of uncultivated marine archaea from comparative metagenomic analyses of bathypelagic plankton of different oceanic regions. *ISME J.* 2:865–886.
- Moreira D, Rodríguez-Valera F, López-García P. 2004. Analysis of a genome fragment of a deep-sea uncultivated Group II euryarchaeote containing 16S rDNA, a spectinomycin-like operon and several energy metabolism genes. *Environ Microbiol.* 6:959–969.
- Nelson-Sathi S, et al. 2012. Acquisition of 1,000 eubacterial genes physiologically transformed a methanogen at the origin of Haloarchaea. *Proc Natl Acad Sci U S A.* 109:20537–20542.
- Nicol GW, Schleper C. 2006. Ammonia-oxidising Crenarchaeota: important players in the nitrogen cycle? *Trends Microbiol.* 14:207–212.
- Ochman H, Lawrence JG, Groisman EA. 2000. Lateral gene transfer and the nature of bacterial innovation. *Nature* 405:299–304.
- Pester M, Schleper C, Wagner M. 2011. The Thaumarchaeota: an emerging view of their phylogeny and ecophysiology. *Curr Opin Microbiol.* 14:300–306.
- Price MN, Dehal PS, Arkin AP. 2009. FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol.* 26:1641–1650.
- Ragon M, Van Driessche AE, García-Ruiz JM, Moreira D, López-García P. 2013. Microbial diversity in the deep-subsurface hydrothermal aquifer feeding the giant gypsum crystal-bearing Naica Mine, Mexico. *Front Microbiol.* 4:37.
- Raychaudhuri S, Stuart JM, Altman RB. 2000. Principal components analysis to summarize microarray experiments: application to sporulation time series. *Pac Symp Biocomput.* 455–466.
- Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European molecular biology open software suite. *Trends Genet.* 16:276–277.
- Rinke C, et al. 2013. Insights into the phylogeny and coding potential of microbial dark matter. *Nature* 499:431–437.
- Saeed AI, et al. 2003. TM4: a free, open-source system for microarray data management and analysis. *Biotechniques* 34:374–378.
- Schleper C, Jurgens G, Jonuscheit M. 2005. Genomic studies of uncultivated archaea. *Nat Rev Microbiol.* 3:479–488.
- Spang A, et al. 2012. The genome of the ammonia-oxidizing *Candidatus Nitrososphaera gargensis*: insights into metabolic versatility and environmental adaptations. *Environ Microbiol.* 14:3122–3145.
- Teeling H, Meyerdieks A, Bauer M, Amann R, Glockner FO. 2004. Application of tetranucleotide frequencies for the assignment of genomic fragments. *Environ Microbiol.* 6:938–947.
- Teeling H, Waldmann J, Lombardot T, Bauer M, Glockner FO. 2004. TETRA: a web-service and a stand-alone program for the analysis and comparison of tetranucleotide usage patterns in DNA sequences. *BMC Bioinformatics* 5:163.
- Treangen TJ, Rocha EP. 2011. Horizontal transfer, not duplication, drives the expansion of protein families in prokaryotes. *PLoS Genet.* 7: e1001284.
- Walker CB, et al. 2010. *Nitrosopumilus maritimus* genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proc Natl Acad Sci U S A.* 107:8818–8823.
- Wolf YI, Makarova KS, Yutin N, Koonin EV. 2012. Updated clusters of orthologous genes for Archaea: a complex ancestor of the Archaea and the byways of horizontal gene transfer. *Biol Direct.* 7:46.

Associate editor: Bill Martin