

- International Ocean-Colour Coordinating Group*. International Ocean-Colour Coordinating Group, Dartmouth, Nova Scotia, 43 pp.
- IOCCG Report Number 3. (2001). Remote sensing of ocean colour in coastal and other optically-complex waters. Sathyendranath, S. (ed) *Report of the International Ocean-Colour Coordinating Group*, International Ocean-Colour Coordinating Group, Dartmouth, Nova Scotia 140 pp.
- Jacquet, S., Partensky, F., Lennon, J., and Vaultot, D. (2001). Diel Patterns of Growth and Division in Marine Picoplankton in Culture. *Journal of Phycology*. 37: 357-369.
- Jaffe, J. S. (2005) Sensing plankton: Acoustics and optical imaging. In: Cullen, J.J. (ed) *Realtime Observation Systems for Ecosystem Dynamics and Harmful Algal Blooms*. UNESCO, Paris, p. (this volume).
- Jeffrey, S. W., Mantoura, R. F. C., and Wright, S. W. (eds). *Phytoplankton Pigments in Oceanography*. UNESCO, Paris, 661 pp.
- Johnsen, G. and Sakshaug, E. (2000). Monitoring of harmful algal blooms along the Norwegian coast using bio-optical methods. *South African Journal of Marine Science*. 22: 309-321.
- Johnson, K. S. and Coletti, L. J. (2002). In situ ultraviolet spectrophotometry for high resolution and long-term monitoring of nitrate, bromide and bisulfide in the ocean. *Deep-Sea Research I*. 49: 1291-1305.
- Kahru, M. and Mitchell, B. G. (2001). Seasonal and nonseasonal variability of satellite-derived chlorophyll and colored dissolved organic matter concentration in the California Current. *Journal of Geophysical Research*. 106: 2517-2529.
- Kaku, M. (1997). *Visions: How Science Will Revolutionize the 21st Century?* Anchor Books Doubleday, New York, 403 pp.
- Kamykowski, D. (1974). Possible interactions between phytoplankton and semidiurnal internal tides. *Journal of Marine Research*. 32: 67-89.
- Keafer, B. A. and Anderson, D. M. (1993). Use of remotely-sensed sea surface temperatures in studies of *Alexandrium tamarense* bloom dynamics. In: Smayda, T.J.; Shimizu, Y (eds) *Toxic Phytoplankton Blooms in the Sea*. Elsevier, Amsterdam, p. 763-768.
- Kerfoot, J. M., Mahoney, K., Kirkpatrick, G., Lohrenz, S., Moline M., and Schofield, O. (2002). Vertical migration of a toxic *Karenia brevis* red tide and the impact on remote sensing reflectance. *EOS Transactions*. AGU/ASLO Ocean Sciences Meeting, Honolulu, HI, 11-15 February.
- Kirk, J. T. O. (1994). *Light and Photosynthesis in Aquatic Ecosystems*. Cambridge University Press, second ed., Cambridge, UK, 509 pp.
- Kirkpatrick, G. J., Millie, D. F., Moline, M. A., and Schofield, O. (2000). Optical discrimination of a phytoplankton species in natural mixed populations. *Limnology and Oceanography*. 45: 467-471.
- Lam, C. W. Y. and Ho, K. C. (1989). Red tides in Tolo Harbour, Hong Kong. In: Nemoto, T. (ed) *Red Tides: Biology, Environmental Science, and Toxicology*. Elsevier, New York, p. 49-52.
- Langlois, G. (2003). Chief, Marine Biotxin Monitoring and Control Program, California Department of Health Services, 2151 Berkeley Way, Berkeley, CA, 94704, pers. comm.