

- Smith, S. L., Pieper, R., Moore, M., Rudstam, L., Green, C., Zamon, J., Flagg, C., and Williamson, C. (1992). Acoustic techniques for in situ observations of zooplankton, In: *Archiv fur Hydrobiologie*, Beiheft 36, Ergebnisse der Limnologie, 35, p. 23-43.
- Sosik, H. M., Olson, R. J., Neubert, M. G., Shalapyonok, A. A., and Solow, A. R. (2003). Growth rates of coastal phytoplankton from time-series measurements with a submersible flow cytometer. *Limnology and Oceanography*. 48: 1756-1765.
- Stramska, M. and Dickey, T. D. (1992). Variability of bio-optical properties in the upper ocean associated with diel cycles in phytoplankton population. *Journal of Geophysical Research*. 97: 17,873-17,887.
- Stramski, D., Bricaud, A., and Morel, A. (2001). Modeling the inherent optical properties of the ocean based on the detailed composition of planktonic community. *Applied Optics*, 40: 2929-2945.
- Strutton, P. G. and Chavez, F. P. (2000). Primary productivity in the equatorial Pacific during the 1997-98 El Niño. *Journal of Geophysical Research*, 105: 26,089-26,101.
- Stumpf, R. P. (2001). Applications of satellite ocean color sensors for monitoring and predicting harmful algal blooms. *Human and Ecological Risk Assessment*. 7: 1363-1368.
- Stumpf, R. P., Culver, M. E., Tester, P. A., Tomlinson, M., Kirkpatrick, G. J., Pederson, B. A., Truby, E., Ransibrahmanakul, V., and Soracco, M. (2003). Monitoring *Karenia brevis* blooms in the Gulf of Mexico using satellite ocean color imagery and other data. *Harmful Algae*. 2: 147-160.
- Tester, P. A. and Steidinger, K. A. (1997). *Gymnodinium breve* red tide blooms: Initiation, transport, and consequences of surface circulation. *Limnology and Oceanography*. 42: 1039-1051.
- Thompson, R. O. R. Y. (1979). Coherence significance levels. *Journal of Atmospheric Science*. 36: 2020– 2021.
- Tokar, J. M. and Dickey, T. D. (2000). Chemical sensor technology – Current and future applications. In: Varney, M. S. (ed) *Chemical Sensors in Oceanography*. Gordon & Breach Scientific Publishers, Amsterdam, p. 303-329.
- Trowbridge, J. H. and Nowell, A. R. M. (1994). An introduction to the Sediment TRansport Events on Shelves and Slopes (STRESS) program. *Continental Shelf Research*. 14: 1057-1061.
- Turner, J. T. and Tester, P. A. (1997). Toxic marine phytoplankton, zooplankton grazers, and pelagic food webs. *Limnology and Oceanography*. 42: 1203-1214.
- Twardowski, M. S., Boss, E., Macdonald, J. B., Pegau, W. S., Barnard, A. H., and Zaneveld, J. R. V. (2001). A model for estimating bulk refractive index from the optical backscattering ratio and the implications for understanding particle composition in case I and case II waters. *Journal of Geophysical Research*. 106: 14,129-14,142.
- Ulloa, O., Sathyendranath, S., and Platt, T. (1994). Effect of the particle size-distribution on the backscattering ratio in seawater. *Applied Optics*. 33: 7070-7077.
- Vaulot, D., and Marie, D. (1999). Diel variability of photosynthetic picoplankton in the equatorial Pacific. *Journal of Geophysical Research*. 104: 3297-3310.
- Vaulot, D., Marie, D., Olson, R. J., and Chisholm, S. W. (1995). Growth of *Prochlorococcus*, a photosynthetic prokaryote, in the equatorial Pacific Ocean. *Science*. 268: 1480-1482.