

- Walsh, J. J. and Steidinger, K. A. (2001). Saharan dust and Florida red tides: The cyanophyte connection. *Journal of Geophysical Research*. 107: 11,597–11,612.
- Widder, E. A., Case, J. F., Bernstein, S. A., MacIntyre, S., Lowenstine, M. R., Bowlby, M. R., and Cook, D. P. (1993). A new large volume bioluminescence bathyphotometer with defined turbulence excitation, *Deep-Sea Research I*, 40: 607-627.
- Yanagi, T. and Okaichi, T. (1997). Seto Inland Sea - historical background. In: Yanagi, T. (ed) *Sustainable Development in the Seto Inland Sea, Japan — From the Viewpoint of Fisheries*. Terra Scientific Publishing Company, Tokyo, p. 9-14.
- Yoder, J. A., Moore, J. K., and Swift, R. N. (2001). Putting together the big picture: remote-sensing observations of ocean color. *Oceanography Magazine*. 14: 33-40.
- Yu, X., Dickey, T., Bellingham, J., Manov, D., and Streitlen, K. (2002). The application of autonomous underwater vehicles for interdisciplinary measurements in Massachusetts and Cape Cod Bays. *Continental Shelf Research*. 22: 2225-2245.
- Zaneveld, J. R. V., Moore, C., Barnard, A. H., Twardowski, M., and Chang, G. C. (2004). Correction and analysis of spectral absorption data taken with the WET Labs ac-s, *Ocean Optics XVII*, Fremantle, Australia, 22-26 October.
- Zingone, A. and Enevoldsen, H. O. (2000). The diversity of harmful algal bloom: A challenge for science and management. *Ocean & Coastal Management*. 43: 725-748.